

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR DISINFECTING: (Bactericidal, Fungicidal, Virucidal)

For use on hard, non-porous non-food contact surfaces such as floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, and plastic surfaces. Remove gross filth and heavy soil deposits then thoroughly wet surfaces. Use 2 oz per gallon of water (8 oz per gallon against Canine parvovirus and Porcine parvovirus) for a minimum contact time of 10 minutes in a single application. Can be applied with a cloth, mop, or sponge, as well as by coarse spray or soaking. The use solution is prepared fresh for each use then discarded. Rinsing is not necessary unless floors are to be waxed or polished.

FUNGICIDAL DIRECTIONS:

For use in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities follow disinfection directions.

MILDEWSTATIC INSTRUCTIONS:

Will effectively control the growth of mold and mildew plus the odors caused by them when applied to hard, nonporous surfaces such as walls, floors, and table tops. Apply solution (2 oz per gallon of water) with a cloth, mop, sponge or coarse spray making sure to wet all surfaces completely. Let air dry. Repeat application weekly or when growth reappears.

VETERINARY CLINICS / ANIMAL CARE / ANIMAL LIFE SCIENCE LABORATORIES / ZOOS / PET SHOPS / KENNELS / BREEDING AND GROOMING ESTABLISHMENTS / EQUINE FARMS DISINFECTION DIRECTIONS:

For disinfecting the following hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a solution of 2 oz of KennelSol per gallon of water (or equivalent dilution) (8 oz per gallon against Canine parvovirus or Porcine parvovirus), using a cloth, mop, sponge, spray, or soaking for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

*FOR CANINE PARVOVIRUS (AND PORCINE PARVOVIRUS) ACTIVITY:

KennelSol is effective against Canine Parvovirus (and Porcine parvovirus) at 8 oz per gallon. Follow Veterinary Practice / Animal Care / Animal Laboratory / Zoos / Pet Shops / Kennels Disinfection Directions.

**SPECIAL INSTRUCTIONS FOR INACTIVATING AVIAN INFLUENZA:

Remove all poultry and feeds from premises, trucks, coops, and crates. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with the disinfecting solution (2 oz per gallon of water) (660 ppm active quat) (equivalent use dilution) for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers used in removing litter and manure. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

KennelSol®

Germicidal Detergent & Deodorant

Disinfects, Cleans and Deodorizes in one labor saving step!



ACTIVE INGREDIENTS:

Didecyl dimethyl ammonium chloride.....	2.535%
n-Alkyl (C ₁₄ 50%, C ₁₂ 40%, C ₁₆ 10%) dimethyl benzyl ammonium chloride.....	1.690%
OTHER INGREDIENTS.....	95.775%
TOTAL.....	100.000%

KEEP OUT OF REACH OF CHILDREN DANGER

(See bottom of label for additional Precautionary Statements, First Aid, and Directions for Use)

Net Contents: 5 Gallons (18.927 litres)

EPA Reg. No. 62472-1 EPA Est. No. 10634-MA-1

For use in: Kennels, pet shops, veterinary clinics, animal life science laboratories, breeding establishments, grooming establishments, schools, colleges, federally inspected meat and poultry plants, equine farms, tack shops, zoos, poultry farms, turkey farms, dairy farms, hog farms, and households. Disinfects, cleans, and deodorizes the following hard nonporous non-food contact inanimate surfaces: floors, walls, non-medical metal surfaces, non-medical stainless steel surfaces, glazed porcelain, plastic surfaces (such as polypropylene, polystyrene, etc.).

Recommended for use in: Federally inspected meat and poultry plants as a disinfectant agent for use in all departments. Food products and packaging materials must be removed from the room or carefully protected. Use product in accordance with its label. All surfaces must be thoroughly rinsed with potable water.

KennelSol® Germicidal Detergent & Deodorant

A multi-purpose germicidal detergent and deodorant effective in hard waters up to 400 ppm hard water (calculated as CaCO₃) plus 5% organic serum. Disinfects, cleans, and deodorizes in one labor saving step.

For use in kennels, pet shops, tack shops, veterinary clinics, animal life science laboratories, breeding & grooming establishments, schools, colleges, equine farms, airports, hotels & motels. For use in households.

Bactericidal against the following pathogenic bacteria according to the AOAC Use-Dilution Test method, current edition, modified in the presence of 400 ppm synthetic hard water (calculated as CaCO₃) plus 5% organic serum:

Proteus vulgaris	Shigella sonnei
Bordetella bronchiseptica	Staphylococcus epidermidis (antibiotic resistant)
Serratia marcescens	Enterobacter cloacae
Escherichia coli	Pasteurella multocida
Staphylococcus aureus (antibiotic resistant)	Staphylococcus aureus
Fusobacterium necrophorum	Salmonella typhi
Klebsiella pneumoniaea (antibiotic resistant)	Chlamydia psittaci
Streptococcus pyogenes	Shigella flexneri
Proteus mirabilis	Enterobacter aerogenes
Salmonella enterica	Streptococcus faecalis (antibiotic resistant)
Salmonella typhimurium	Klebsiella pneumoniae
Escherichia coli (antibiotic resistant)	Streptococcus faecalis

Fungicidal against Trichophyton mentagrophytes and Candida albicans according to the AOAC fungicidal test, modified in the presence of 400 ppm hard water (calculated as CaCO₃) plus 5% organic serum.

Virucidal against *Canine parvovirus, Canine distemper, Feline leukemia, Feline panleukopenia, Feline picornavirus, Influenza A/Hong Kong, Herpes simplex type 1, Herpes simplex type 2, Vaccinia, Rubella, Adenovirus type 4, Rabies, Porcine parvovirus, Pseudorabies, Infectious bovine rhinotracheitis, Infectious bronchitis (Avian IBV), **Avian Influenza A (H3N2) virus (Avian Reassortant) (ATCC VR-2072), and **Avian Influenza A (H5N1) virus according to the virucidal qualification, modified in the presence of 400 ppm hard water (calculated as CaCO₃) plus 5% organic serum.



Manufactured for:

Alpha Tech Pet, Inc.

25 Porter Road, Suite 210
Littleton, MA 01460 • 800-222-5537

© 2019 Alpha Tech Pet Incorporated • Made and Printed in U.S.A.



STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container in areas inaccessible to children. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your local State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Fill container 1/4 full

with water and recap. Shake 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

*PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. Wear protective eyewear (goggles, face shield or safety glasses), protective clothing and protective gloves (rubber or chemical resistant). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. This pesticide is toxic to fish, aquatic invertebrates, oysters, and shrimp. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.